

CARVE: L4 Gridded Footprints from WRF-STILT model, 2012-2016

Get Data

Documentation Revision Date: 2022-05-06

Dataset Version: 1.1

Summary

This data set provides Weather Research and Forecasting (WRF) Stochastic Time-Inverted Lagrangian Transport (STILT) Footprint data products for particle receptors located at positions along Carbon in Arctic Reservoirs Vulnerability Experiment (CARVE) flight paths (2012 - 2015) and various meteorological stations in Alaska and the Canadian Arctic. Each product consists of multiple NetCDF footprint files packaged as a TAR/GZIP file. These aircraft and station positions were treated as receptors in the WRF-STILT model in order to simulate the land surface influence on observed atmospheric constituents. The measurements included in this data set are crucial for understanding changes in Arctic carbon cycling and the potential threats posed by thawing of Arctic permafrost.

There are 87 TAR/GZIP files included in this data set containing a total of 666,433 WRF-STILT Footprint files in NetCDF format.




Figure 1. Surface contours (50% --- blue; 80% --- purple) for the average WRF-STILT influence functions (mid-afternoon averages 2012-2014) show CO₂ and CH₄ measurements from the CARVE tower have a high sensitivity to the boreal forests and lowlands of interior Alaska. Cyan and red circles are shown for the footprints after filtering (from Karion et al., 2016).

Citation

Henderson, J., J.B. Miller, T. Nehrkorn, R.Y-W. Chang, C. Sweeney, N. Steiner, S.C. Wofsy, and C.E. Miller. 2017. CARVE: L4 Gridded Footprints from WRF-STILT model, 2012-2016. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAC/1431>

Table of Contents

1. [Dataset Overview](#)

2. [Data Characteristics](#)
3. [Application and Derivation](#)
4. [Quality Assessment](#)
5. [Data Acquisition, Materials, and Methods](#)
6. [Data Access](#)
7. [References](#)
8. [Dataset Revisions](#)

1. Dataset Overview

Project: Carbon in Arctic Reservoirs Vulnerability Experiment ([CARVE](#))

The Carbon in Arctic Reservoirs Vulnerability Experiment (CARVE) is a NASA Earth Ventures (EV-1) investigation designed to quantify correlations between atmospheric and surface state variables for Alaskan terrestrial ecosystems through intensive seasonal aircraft campaigns, ground-based observations, and analysis sustained over a 5-year mission. CARVE collected detailed measurements of greenhouse gases on local to regional scales in the Alaskan Arctic and demonstrated new remote sensing and improved modeling capabilities to quantify Arctic carbon fluxes and carbon cycle-climate processes.

Related Data:

These modeled footprint data were based on input from the [CARVE: L4 Gridded Particle Trajectories for WRF-STILT model, 2012-2016](#) data available from ORNL DAAC. The relationship between particle files and footprint files is outlined in the companion file 'carve_wrf-stilt_inventory.csv'.

A full list of CARVE data products is available at: <https://carve.ornl.gov/dataproducts.html>

2. Data Characteristics

Spatial Coverage: Campaign flights and station observations within the CARVE domain

Spatial Resolution: 0.25-degree for *foot1* data; 0.1-degree for *footnearfield1* data

Temporal Coverage: 20120523 - 20160428

Temporal Resolution: Hourly

Study Area (coordinates in decimal degrees)

Site	Westernmost Longitude	Easternmost Longitude	Northernmost Latitude	Southernmost Latitude
CARVE Domain	-180	180	90	30

Data File Information

The TAR/GZIP files (*.tar.gz) contain NetCDF files representing gridded footprints from WRF-STILT simulations for one particle receptor location. Each file aggregates particle footprints on a lat/lon/time grid starting at the STILT simulation start time.

Notes about TAR/GZIP files

- If *convect* is present in the filename, STILT was run with convective fluxes active. If *noconvect* is present, or if there is no label, then fluxes are not active (standard for simulations through 2015).
- Simulations for 2012 were run with WRF v3.4.1. Simulations for 2013 were run with both v3.4.1 and v3.5.1. Subsequent years used only v3.5.1.
- *CARVE-CAN* refers to use of the WRF *CARVE-CAN* domain that was available after 2014.
- Filenames containing *freum** were generated going back 15 days instead of the usual 10.

The first surface influence field, represented by the *foot1* variable in the NetCDF files, provides 10 days of surface influence representing the response of the receptor to a unit surface emission (ppm/nmol m⁻² s⁻¹) of CO₂ in each 0.25- x 0.25-degree grid cell within the whole area of coverage (30N to 90N, 180E to 180W) at hourly temporal resolution.

The second surface influence field, represented by the *footnearfield1* variable in the NetCDF files, provides 24 hours of surface influence representing the response of the receptor to a unit surface emission (ppm/nmol m⁻² s⁻¹) of CO₂ in each 0.1- x 0.1-degree grid cell within a small region close to the measurement location at hourly temporal resolution.

The WRF-STILT model as run for this project is described further in Henderson et al. (2015). File naming convention for WRF-STILT footprint files is described in Table 1.

Table 1. Naming convention for NetCDF files generated by WRF-STILT.

Example file name: *foot2013x06x25x04x00x65.1330Nx147.4539Wx00003.nc*

Name element	Example value	Units
Year	2013	YYYY
Month	06	MM
Day	25	DD
Hour	04	hh (UTC)
Minute	00	mm (UTC)
Latitude	65.1330N	decimal degrees

Longitude	127.4539W	decimal degrees
Height A.G.L.	00003	m

For example, the above file contains the modeled particle footprints for June 25, 2013 at 4:00 UTC. The observation was taken at 65.1330N, 127.4539W at 3 m above ground level.

Data Variables

Table 2. Data variables in each NetCDF file. Fill values or missing data were set to -1.0E34 for all variables.

Variable name	Units	Description
foot1	ppm per (umol m-2 s-1)	gridded STILT footprint
foot1date	days	days since 2000-01-01 00:00:00 UTC
foot1hr	hours	hours back from STILT start time
foot1lat	degrees_north	degrees latitude of center of grid cells
foot1lon	degrees_east	degrees longitude of center of grid cells
footnearfield1	ppm per (umol m-2 s-1)	gridded STILT footprint
footnearfield1date	days	days since 2000-01-01 00:00:00 UTC
footnearfield1hr	hours	hours back from STILT start time
footnearfield1lat	degrees_north	degrees latitude of center of grid cells
footnearfield1lon	degrees_east	degrees longitude of center of grid cells
nchar		identifier string

3. Application and Derivation

WRF-STILT footprints support accurate estimates of CO₂ and CH₄ surface-atmosphere fluxes using CARVE flight and tower observations. Simulated CO₂ mole fractions from the PolarVPRM model based on WRF-STILT footprints show strong agreement with tower observations, suggesting that the WRF-STILT model does a good job representing the meteorology of the region (Karion et al., 2016).

4. Quality Assessment

Preliminary analysis in support of CARVE showed good overall agreement between WRF outputs and quality-controlled surface and radiosonde observations.

Analysis of STILT footprints showed realistic seasonal variability and good agreement with tower observations, indicating that WRF-STILT footprints are of high quality and support accurate estimates of CO₂ and CH₄ surface-atmosphere fluxes using CARVE observations (Henderson et al. 2015).

5. Data Acquisition, Materials, and Methods

CARVE Flights

These data represent one part of the data collected by the Carbon in Arctic Reservoirs Vulnerability Experiment (Miller et al, 2012). A C-23 Sherpa aircraft made frequent flights out of Fairbanks, Alaska between March and November over multiple years, observing the spring thaw, summer draw-down, and fall refreeze of the Arctic growing season. Flights concentrate observations on three study domains: the North Slope, the interior, and the Yukon River valley. North Slope flights cover regions of tundra and continuous permafrost and were anchored by flux towers in Barrow, Atkasuk, and Igotuk. Flights to Prudhoe Bay characterize the CO₂ and CH₄ emissions from oil and natural gas processing plants. Flights over interior Alaska sample discontinuous permafrost, boreal forests, and wetlands. A complete list of CARVE flights can be found at: <https://carve.ornl.gov/flights.html>. Flight paths and atmospheric gas concentrations for CARVE surveys can be visualized through the CARVE Flight Data Visualization Tool (<http://carve.ornl.gov/visualize>).




Figure 2. CARVE flights during 2012-2015 delivered measurements over continuous and discontinuous permafrost regimes.

The CARVE aircraft carried a remote sensing and atmospheric sampling payload consisting of the following instruments: a Fourier transform spectrometer (FTS), and an in situ gas analyzer suite (ISGAS) with a gas analyzer and flask sampling system (see <https://carve.ornl.gov/documentation.html>). All instruments were controlled by a master computer system (Data Acquisition and Distribution System, DADS). DADS also recorded GPS data (Lat, Lon, altitude), aircraft pitch, roll, and yaw, as well as basic meteorological data from onboard instruments.

WRF-STILT Simulations

CARVE aircraft positions and various flux tower locations were treated as receptors in a Stochastic Time-Inverted Lagrangian Transport (STILT) model coupled with meteorology fields from the polar variant of the Weather and Research Forecasting (WRF; Skamarock et al., 2008) model, in order to model the land surface influence on observed atmospheric constituents. The atmospheric model was configured to generate high-quality, high-resolution meteorological fields over Arctic and boreal Alaska.

The WRF-STILT coupled model is described in Nehr Korn et al. (2010).

STILT applies a Lagrangian particle dispersion model backwards in time from a measurement location (the "receptor" location), to create the adjoint of the transport model in the form of a "footprint" field (Nehr Korn et al., 2010; Henderson et al., 2015). The footprint, with units of mixing ratio (ppm --- CO₂; ppb -- - CH₄) per (umol m⁻² s⁻¹), quantifies the influence of upwind surface fluxes on concentrations measured at the receptor and is computed by counting the number of particles in a surface-influenced volume and the time spent in that volume. Gridded footprints are provided as NetCDF files.

6. Data Access

These data are available through the Oak Ridge National Laboratory (ORNL) Distributed Active Archive Center (DAAC).

[CARVE: L4 Gridded Footprints from WRF-STILT model, 2012-2016](#)

Contact for Data Center Access Information:

- E-mail: uso@daac.ornl.gov
- Telephone: +1 (865) 241-3952

7. References

- Henderson, J.M., J. Eluszkiewicz, M.E. Mountain, T. Nehr Korn, R.Y.-W. Chang, A. Karion, J.B. Miller, C. Sweeney, N. Steiner, S.C. Wofsy, and C.E. Miller. 2015. Atmospheric transport simulations in support of the Carbon in Arctic Reservoirs Vulnerability Experiment (CARVE). *Atmos. Chem. Phys.* 15:4093-4116.
- Karion, A., C. Sweeney, J.B. Miller, A.E. Andrews, R. Commane, S. Dinardo, J.M. Henderson, J. Lindaas, J.C. Lin, K.A. Luus, T. Newberger, P. Tans, S.C. Wofsy, S. Wolter, and C.E. Miller. 2016. Investigating Alaskan methane and carbon dioxide fluxes using measurements from the CARVE tower. *Atmos. Chem. Phys.* 16:5383-5398.
- Miller, C.E., Dinardo, S.J. et al. (2012). CARVE: The Carbon in Arctic Reservoirs Vulnerability Experiment. , *2012 IEEE Aerospace Conference*. <http://dx.doi.org/10.1109/AERO.2012.6187026>
- Nehr Korn, T., J. Eluszkiewicz, S.C. Wofsy, J.C. Lin, C. Gerbig, M. Longo, and S. Freitas. 2010. Coupled weather research and forecasting-stochastic time-inverted lagrangian transport (WRF-STILT) model. *Meteorol. Atmos. Phys.* 107:51-64. doi:10.1007/s00703-010-0068-x
- Skamarock, W.C. and J.B. Klemp. 2008. A time-split nonhydrostatic atmospheric model for weather research and forecasting applications. *Journal of Computational Physics*, 227(7): 3465-3485.

8. Dataset Revisions

This dataset was revised on May 6, 2022 to rename 10 data files that contained the plus sign in their filenames. The plus sign in each file was replaced with an underscore. No data values in the files were changed.



Home

About Us

- Mission
- Data Use and Citation Policy
- User Working Group
- Partners

Get Data

- Science Themes
- NASA Projects
- All Datasets

Submit Data

- Submit Data Form
- Data Scope and Acceptance
- Data Authorship Policy
- Data Publication Timeline
- Detailed Submission Guidelines

Tools

- MODIS
- THREDDS
- SDAT
- Daymet
- Airborne Data Visualizer
- Soil Moisture Visualizer
- Land - Water Checker

Resources

- Learning
- Data Management
- News
- Earthdata Forum [↗](#)

Contact Us